

SYSTEMATIC LIST OF THE HOVERFLIES OF BEDDINGTON FARMLANDS

This list follows S.Ball and R.Morris 2013, Britain's Hoverflies, Second Edition, Wild Guides. Where species not covered in Ball and Morris systematic order follows Plant (2011) and Collins and Coleman (2009) .

Common Name	Scientific Name	Conservation Status	Beddington Farmlands Status
			Unless otherwise stated records refer to specimens caught in malaise traps by C.Plant (2011) and G.Collins & D.A Coleman (2012). Other records by P.R.Alfrey (PRA) or F.Prater (FP).
	<i>Baccha elongata</i>		Seen along permitted footpath
	<i>Melanostoma mellinum</i>		
	<i>Melanostoma scalare</i>		
	<i>Platycheirus albimanus</i>		Abundant
	<i>Platycheirus aurolateralis</i>		
	<i>Platycheirus angustatus</i>		
	<i>Platycheirus clypeatus</i>		
	<i>Platycheirus fulviventris</i>		
	<i>Platycheirus rosarum</i>		
	<i>Paragus haemorrhous</i>		
	<i>Chrysotoxum bicinctum</i>		
	<i>Chrysotoxum festivum</i>		
	<i>Chrysotoxum cautum</i>		Recorded by PRA 270614
	<i>Chrysotoxum verralli</i>		
	<i>Xanthogramma pedissequum</i>		Recorded regularly by PRA
	<i>Sphaerophoria interrupta</i>		
	<i>Sphaerophoria rueppelli</i>		
	<i>Sphaerophoria scripta</i>		Common
	<i>Leucozona lucorum</i>		Recorded by PRA e.g. 150515

	<i>Dasysyrphus tricinctus</i>		
	<i>Eupeodes luniger</i>		
	<i>Eupeodes corollae</i>		
	<i>Eupoedes latifasciatus</i>		
	<i>Melangyna cincta</i>		
Marmalade Hoverfly	<i>Episyrphus balteatus</i>		Abundant
	<i>Epistrophe eligans</i>		
Common Hoverfly	<i>Syrphus Ribesii</i>		Common
	<i>Syrphus vitripennis</i>		
	<i>Cheilosia pagana</i>		
	<i>Cheilosia vernalis</i>		
	<i>Cheilosia bergenstammi</i>		
	<i>Cheilosia impressa</i>		
	<i>Cheilosia latifrons</i>		
	<i>Cheilosia proxima</i>		
	<i>Rhingia campestris</i>		Seen by PRA e.g. 100613
	<i>Neoascia interrupta</i>	Local	
	<i>Neoascia podagrica</i>		
	<i>Neoascia podagrica</i>		
	<i>Meoascia meticulosa</i>		
	<i>Orthonevra brevicornis</i>		
	<i>Eristalis pertinax</i>		Common
	<i>Eristalis arbustorum</i>		Fairly common
	<i>Eristalis intricaria</i>		Fairly prominent
	<i>Eristalis tenax</i>		Recorded commonly by PRA and FP
	<i>Eristalinus sepulchralis</i>		Regular on 100 acre in rank vegetation.
	<i>Myathropha florea</i>		Recorded commonly by PRA and FP.
	<i>Anasimyia lineata</i>		
	<i>Anasimyia contracta</i>		
	<i>Helophilus hybridus</i>		
	<i>Helophilus pendulus</i>		
	<i>Helophilus trivittatus</i>		Recorded by FP.
	<i>Parhelophilus frutetorum</i>		
	<i>Parhelophilus versicolor</i>		
	<i>Eumerus strigatus</i>		
	<i>Merodon equestris</i>		
	<i>Pipiza noctiluca</i>		
	<i>Pipizella viduata</i>		
	<i>Volucella bombylans</i>		

	<i>Volucella pellucens</i>			
	<i>Volucella zonaria</i>		Recorded commonly by PRA and FP	
	<i>Volucella inanis</i>		One record: 110710 (PRA)	
	<i>Syrirta pipiens</i>			
	<i>Xylota segnis</i>		Recorded by PRA e.g. 120515	
			63 species	

References

S.Ball and R.Morris 2013, Britian's Hoverflies, Second Edition, Wild Guides

C.Plant 2011, Invertebrate Survey, Beddington Farmlands

G.Collins and D.A Coleman, 2009, Malaise Trap at Beddington, Beddington Farmlands Bird and Willdife Report 2012